AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A method of treating metastasis of cancer in a subject at a site distal to the site of surgical resection of a tumor in the subject, wherein said subject is in need of such treatment and the metastasis is present at a site distal to the site of surgical resection of the tumor in the subject, said method comprising the steps of:

surgically resecting a tumor from the subject; and administering an attenuated, replication-competent, oncolytic herpes virus to the site of surgical resection.

- 2. (Canceled).
- 3. (Original) The method of claim 1, wherein said cancer has metastasized from the site of surgical resection.
- 4. (Original) The method of claim 3, wherein said cancer is present in the lymphatic system of said subject.
- 5. (Original) The method of claim 4, wherein said cancer is present in a lymph node of said subject.
- 6. (Original) The method of claim 1, wherein said herpes virus is a herpes simplex-1-derived virus.

- 7. (Original) The method of claim 6, wherein said herpes virus is NV1023.
- 8. (Original) The method of claim 1, wherein said subject is a human.
- 9. (Original) The method of claim 1, wherein said herpes virus is administered to said subject by injection.
- 10. (Original) The method of claim 1, wherein said herpes virus comprises a heterologous nucleic acid molecule encoding a therapeutic product.
- 11. (Original) The method of 10, wherein said therapeutic product is selected from the group consisting of cytotoxins, immunomodulatory proteins, tumor antigens, antisense nucleic acid molecules, and ribozymes.
- 12. (Original) The method of claim 1, further comprising administering a second anticancer treatment to said subject.
- 13. (Original) The method of claim 12, wherein said second anticancer treatment is selected from the group consisting of chemotherapy, biological therapy, radiation therapy, and gene therapy.
 - 14. (Canceled).

15-27. (Canceled).

28. (Currently Amended) A method of treating regional lymphatic metastases in a subject in need of such treatment, wherein the regional lymphatic metastases are present at a site distal to a site of surgical resection of a tumor in the subject, said method comprising the steps of:

surgically resecting a tumor from the subject; and

administering an attenuated, replication-competent, oncolytic herpes virus to the site of surgical resection.